

A FACT SHEET FOR PARENTS

What is a concussion?

A concussion is a brain injury. Concussions are caused by a bump, blow, or jolt to the head or body. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

What are the signs and symptoms?

You can't see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days after the injury. If your teen reports **one or more** symptoms of concussion listed below, or if you notice the symptoms yourself, keep your teen out of play and seek medical attention right away.

Signs Observed	Symptoms Reported
by Parents or Guardians	by Athlete
 Appears dazed or stunned Is confused about	 Headache or "pressure"
assignment or position Forgets an instruction Is unsure of game, score,	in head Nausea or vomiting Balance problems or
or opponent Moves clumsily Answers questions slowly Loses consciousness	dizziness Double or blurry vision Sensitivity to light
(even briefly) Shows mood, behavior,	or noise Feeling sluggish, hazy,
or personality changes Can't recall events prior	foggy, or groggy Concentration or memory
to hit or fall Can't recall events after	problems Confusion Just not "feeling right"
hit or fall	or is "feeling down"

How can you help your teen prevent a concussion?

Every sport is different, but there are steps your teens can take to protect themselves from concussion and other injuries.

• Make sure they wear the right protective equipment for their activity. It should fit properly, be well maintained, and be worn consistently and correctly.

- Ensure that they follow their coaches' rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.

What should you do if you think your teen has a concussion?

- 1. Keep your teen out of play. If your teen has a concussion, her/his brain needs time to heal. Don't let your teen return to play the day of the injury and until a health care professional, experienced in evaluating for concussion, says your teen is symptom-free and it's OK to return to play. A repeat concussion that occurs before the brain recovers from the first—usually within a short period of time (hours, days, or weeks)—can slow recovery or increase the likelihood of having long-term problems. In rare cases, repeat concussions can result in edema (brain swelling), permanent brain damage, and even death.
- **2. Seek medical attention right away.** A health care professional experienced in evaluating for concussion will be able to decide how serious the concussion is and when it is safe for your teen to return to sports.
- **3. Teach your teen that it's not smart to play with a concussion.** Rest is key after a concussion. Sometimes athletes wrongly believe that it shows strength and courage to play injured. Discourage others from pressuring injured athletes to play. Don't let your teen convince you that s/he's "just fine."
- 4. Tell all of your teen's coaches and the student's school nurse about ANY concussion. Coaches, school nurses, and other school staff should know if your teen has ever had a concussion. Your teen may need to limit activities while s/he is recovering from a concussion. Things such as studying, driving, working on a computer, playing video games, or exercising may cause concussion symptoms to reappear or get worse. Talk to your health care professional, as well as your teen's coaches, school nurse, and teachers. If needed, they can help adjust your teen's school activities during her/his recovery.

If you think your teen has a concussion: Don't assess it yourself. Take him/her out of play. Seek the advice of a health care professional.

It's better to miss one game than the whole season.

For more information and to order additional materials *free-of-charge*, visit: www.cdc.gov/Concussion.



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES CENTERS FOR DISEASE CONTROL AND PREVENTION